

---

**Visualization of macrophage recruitment in head and neck carcinoma model using fluorine-19 magnetic resonance imaging.**

**Journal:** Magn Reson Med

**Publication Year:** 2018

**Authors:** Aman Khurana, Fanny Chapelin, Hongyan Xu, Joseph R Acevedo, Alfred Molinolo, Quyen Nguyen, Eric T Ahrens

**PubMed link:** 28748562

**Funding Grants:** Molecular Imaging for Stem Cell Science and Clinical Application

**Public Summary:**

To evaluate the role of infiltrating macrophages in murine models of single and double mutation head and neck tumors using a novel fluorine-19 (19F) MRI technology.

**Scientific Abstract:**

**PURPOSE:** To evaluate the role of infiltrating macrophages in murine models of single and double mutation head and neck tumors using a novel fluorine-19 (19F) MRI technology. **METHODS:** Tumor cell lines single-hit/SCC4 or double-hit/Cal27, with mutations of TP53 and TP53 & FHIT, respectively, were injected bilaterally into the flanks of (n = 10) female mice. With tumors established, perfluorocarbon nanoemulsion was injected intravenously, which labels in situ predominantly monocytes and macrophages. Longitudinal spin density-weighted (19F) MRI data enabled quantification of the macrophage burden in tumor and surrounding tissue. **RESULTS:** The average number of (19F) F atoms within the tumors was twice as high in the Cal27 group compared with SCC4 ( $3.9 \times 10^{19}$ ) and  $2.0 \times 10^{19}$  (19F) F/tumor, respectively;  $P = 0.0034$ ) two days after contrast injection, signifying increased tumor-associated macrophages in double-hit tumors. The difference was still significant 10 days after injection. Histology stains correlated with in vivo results, exhibiting numerous perfluorocarbon-labeled macrophages in double-hit tumors and to a lesser extent in single-hit tumors. **CONCLUSIONS:** This study helps to establish (19F) F MRI as a method for quantifying immune cells in the tumor microenvironment, allowing distinction between double and single-hit head and neck tumors. This technique would be extremely valuable in the clinic for pretreatment planning, prognostics, and post-treatment surveillance. Magn Reson Med 79:1972-1980, 2018. (c) 2017 International Society for Magnetic Resonance in Medicine.

---

**Source URL:** <https://www.cirm.ca.gov/about-cirm/publications/visualization-macrophage-recruitment-head-and-neck-carcinoma-model-using>